

# Habitat simplification increases the impact of a freshwater invasive fish

M. E. Alexander · H. Kaiser · O. L. F. Weyl ·  
J. T. A. Dick

Received: 9 December 2013 / Accepted: 29 April 2014 / Published online: 11 May 2014  
© Springer Science+Business Media Dordrecht 2014

**Abstract** Biodiversity continues to decline at a range of spatial scales and there is an urgent requirement to understand how multiple drivers interact in causing such declines. Further, we require methodologies that can facilitate predictions of the effects of such drivers in the future. Habitat degradation and biological invasions are two of the most important threats to biodiversity and here we investigate their combined effects, both in terms of understanding and predicting impacts on native species. The predatory largemouth bass *Micropterus salmoides* is one of the World's Worst Invaders, causing declines in native prey species, and its introduction often coincides

with habitat simplification. We investigated the predatory functional response, as a measure of ecological impact, of juvenile largemouth bass in artificial vegetation over a range of habitat complexities (high, intermediate, low and zero). Prey, the female guppy *Poecilia reticulata*, were representative of native fish. As habitats became less complex, significantly more prey were consumed, since, even although attack rates declined, reduced handling times resulted in higher maximum feeding rates by bass. At all levels of habitat complexity, bass exhibited potentially population destabilising Type II functional responses, with no emergence of more stabilising Type III functional responses as often occurs in predator-prey relationships in complex habitats. Thus, habitat degradation and simplification potentially exacerbate the impact of this invasive species, but even highly complex habitats may ultimately not protect native species. The utilisation of functional responses under varying environmental contexts provides a method for the understanding and prediction of invasive species impacts.

---

M. E. Alexander (✉)  
Centre for Invasion Biology, Department of Botany and  
Zoology, Stellenbosch University, Matieland 7602, South  
Africa  
e-mail: malexander@sun.ac.za

H. Kaiser  
Department of Ichthyology and Fisheries Science, Rhodes  
University, P.O. Box 94, Grahamstown 6140, South Africa

O. L. F. Weyl  
South African Institute for Aquatic Biodiversity (SAIAB),  
Private Bag 1015, Grahamstown 6140, South Africa

O. L. F. Weyl  
Centre for Invasion Biology, SAIAB, Private Bag 1015,  
Grahamstown 6140, South Africa

J. T. A. Dick  
Institute for Global Food Security, School of Biological  
Sciences, Queen's University Belfast, MBC, 97 Lisburn  
Road, Belfast BT9 7BL Northern Ireland, UK

**Keywords** Habitat complexity · Invasive species ·  
Functional response · Impact · Global change ·  
Freshwater fish

## Introduction

Biodiversity at global, regional and local scales is declining and the conservation of natural resources is under threat from a number of drivers of global change

(Sala et al. 2000; Thomas et al. 2004; Mokany et al. 2012). Such processes are of particular concern in freshwater systems, with drivers such as habitat alteration and invasions by non-native species identified as significant stressors (Saunders et al. 2002; Dudgeon et al. 2006). The modification of freshwater habitats results from a range of human-mediated processes including changes to flow regimens (Bunn and Arthington 2002), dam construction (Nilsson and Berggren 2000), and destruction of vegetation (Radomski and Goeman 2001). Similarly, biological invasions are increasing due to a wide range of human-mediated vectors and pathways (Levine and D'Antonio 2003). In freshwaters, invasive species modify ecosystems through a range of processes including competitive exclusion (Rowles and O'Dowd 2006) and predation (Griffen and Delaney 2007), which may result in dramatic changes to native communities (Crooks 1998).

One of the main challenges in ecology is the ability to predict how drivers of global change, such as biological invasions, may impact ecosystems (Parker et al. 1999; Ricciardi et al. 2013; Dick et al. 2014). Furthermore, research often considers and examines these drivers as independent, single entities (Fazey et al. 2005), however, there is an emerging realisation that biodiversity loss will be better understood and predicted when the relative roles of the major drivers are considered in combination (Facon et al. 2006). It is also recognised that drivers may act synergistically through a variety of pathways (Didham et al. 2007). For example, changes in the traits of an invasive species with habitat alterations, such as predatory efficiency, might result in changes in *per capita* effects with potential wide-ranging consequences for native species population dynamics (Parker et al. 1999).

In aquatic systems, the role of habitat structure, such as that provided by algae and macrophytic plants, is well documented in inter- and intraspecific interactions from a wide range of taxa (Boström et al. 1999; Saha et al. 2009; Gosnell et al. 2012). For a number of fish species in particular, habitat structure has been shown to mediate impacts of fish predation by, for example, creating a physical barrier to predator movement (Savino and Stein 1982). Mediatory effects may also occur in such interactions due to the provision of refuge space for prey (Persson and Eklöv 1995; Beukers and Jones 1997; Anderson 2001; Almany 2004a). Therefore, the loss of structural complexity resulting from habitat degradation may reduce prey survival due to increased predation

vulnerability (Nelson and Bonsdorff 1990). However, predators that adopt a sit-and-wait strategy of prey capture may perform less efficiently with degradation of habitat, reducing predation success (Flynn and Ritz 1999).

A promising methodology that not only provides an understanding of predator-prey interactions but allows predictions of invasive species impact is to examine the functional response (Dick et al. 2013, 2014; Alexander et al. 2014), that is, the relationship between prey density and predator consumption rate (Solomon 1949; Holling 1959). Such a focus allows important density-dependent effects of predation on population stability to be examined owing to the different contributions of response Types (I, II or III) to population dynamics (Murdoch and Oaten 1975). In a Type I response, predator consumption increases linearly with prey number until a threshold density plateau is reached. However, under certain ranges of prey density, a Type II inversely density-dependent response can result in an increase in mortality risk to prey with decreasing density (Hassell 1978). This is in contrast to reduction in mortality risk when prey density declines in a Type III response (Hassell 1978). This is particularly important in habitat complexity studies, where changes in structure can result in alterations to the functional response Type and hence prey population viability (Lipcius and Hines 1986; Buckel and Stoner 2000; Alexander et al. 2012). Furthermore, the application of functional responses in invasion biology has been demonstrated to be effective, with higher functional responses of invasive species compared to natives in laboratory studies corroborated by results from field studies (Bollache et al. 2008; Dick et al. 2013). Here, we use functional responses to predict the impact that changes in habitat complexity, representative of those resulting from habitat destruction, may have on the predator-prey dynamics of an invasive fish predator, one of the “World’s Worst Invaders” (ISSG 2013), on prey species.

As a result of its popularity as an angling species, the largemouth bass *Micropterus salmoides* is one of the five globally most introduced fish species (Welcomme 1992) and, where such introductions have occurred, predation by this species has a major impact on fish communities (Godinho and Ferreira 2000; Ellender et al. 2011; Almeida et al. 2012). In South Africa and in the Iberian Peninsula in southern Europe, for example, largemouth bass are a well-established invasive species that have subsequently invaded a number of

headwater streams, where many native fish species are now endangered or absent (Ellender et al. 2011; Almeida et al. 2012). In addition to this, such systems are also facing a double threat of habitat degradation due to destruction of natural vegetation (Saunders et al. 2002). As ambush predators, largemouth bass typically use structural littoral habitats including aquatic vegetation (Savino and Stein 1989a), and juveniles in particular predominantly select such environments (Olson et al. 2003). We therefore manipulated simulated-habitat complexity along a prey density gradient in order to ask questions regarding the density-dependence of impact of this invasive predator on prey population in the context of habitat degradation.

The aim of this study was thus to describe and quantify the functional responses, and hence impacts, of juvenile largemouth bass on a prey species, the guppy *Poecilia reticulata*, that served as a commercially available surrogate for endangered fish found in headwater systems, with respect to varying levels of habitat complexity. The aims were to establish whether: (1) juvenile largemouth bass exhibit predatory functional responses towards small fish prey; (2) the functional response Type is habitat dependent; and (3) varying habitat complexity alters the strength of the functional response.

## Materials and methods

### Collection and maintenance of experimental animals

Juvenile largemouth bass *Micropterus salmoides* were collected in June 2013 by electrofishing from Douglas Dam (33°19'16"S; 26°31'15" E) and Grey Dam (33°19'29"S; 26°31'39" E), Grahamstown, South Africa. All fish were transported to the Department of Ichthyology and Fisheries Science (DIFS), Rhodes University, Grahamstown and were housed in 600 L tanks in a closed recirculating system. Fish were allowed to acclimate for at least one week prior to use in predation trials and were maintained on a diet of earthworms. As earthworms were not the focal prey species used in the experiment, this diet ensured that no prior prey learning occurred in holding tanks. Prey used were females of the guppy *Poecilia reticulata* (15–20 mm total length), sourced from a captive breeding stock at the DIFS. Female guppies were selected over males owing to their inconspicuous colouration. Guppies were housed in two

600 L tanks and were fed daily on commercially available fish food.

### Experimental set up

Functional response experiments were conducted in 15 square (1,000 mm) 300 L fibreglass tanks that were part of the same flow-through system as the holding tanks described above (water flow 1 L min<sup>-1</sup>; 23.01±0.17°C, mean temperature ± SE). Tank depth was 370 mm with water level at 300 mm at the central outflow pipe (see Jones 2003 for further detail). In the centre of each tank there was an outlet for water overflow that was covered with mesh and secured with cable ties to prevent predators and/or prey escaping. To reduce potential stress on the fish, each tank was half covered with a dark screen to provide a darkened refuge. An airline provided further aeration of water in the tank in addition to the aerated inflowing water from the recirculating system during predation trials.

To simulate habitat structure, strips of green polyethylene (40 mm long and 15 mm wide) were tied in a uniform arrangement to green plastic mesh, cut to fit the bottom of the aquarium. The mesh was then weighted to the bottom of the tanks. This allowed the artificial vegetation to float upwards and occupy the entire water column in the same way as rooted aquatic plants in freshwater systems (personal observation). A plant mimic was used here to allow for standardisation of cover. Densities of vegetation represented high (2,700 blades m<sup>-2</sup>), medium (1,800 blades m<sup>-2</sup>), low (900 blades m<sup>-2</sup>) and zero habitat complexities. To control for the presence of mesh contributing a further element of habitat complexity, mesh without artificial plants was added in zero complexity treatments.

Bass ( $n=18$ ) were selected from a relatively uniform size class to reduce the influence of size-related differences in prey consumption (86.86±2.49 mm, mean total length ± SE; 14.0±0.38 mm, mean gape height ± SE). Gape height was measured by opening the mouth to its maximum capability without distortion or stretching of the jaws and taking a measurement using callipers. Bass were reused in the four habitat treatments (detailed below), however, we ensured that each individual predator was used a maximum of four times and only once within each prey density in each habitat complexity. At least two days recovery time was allowed between uses.

## Functional response trials

Bass were randomly selected from their holding tanks 24 h prior to a trial and transferred to an experimental tank, where they were held without food to allow for acclimatisation and standardisation of hunger levels. Individual fish were then presented with guppies at six prey densities (2, 4, 8, 16, 32, 64), with at least three replicates per density. Feeding trials were initiated at 10:00 h and prey consumption was examined after four hours. Controls were three replicates of each prey density in the absence of predators at each of the habitat complexities.

## Statistical analysis

All analysis was carried out in R v. 2.15.1 (R Development Core Team 2012). Differences in overall prey consumption among habitat complexities and prey densities were assessed using a generalised linear model (GLM) with binomial error distribution. As no interaction was found between habitat complexity and prey density, the interaction term was removed to identify the minimum adequate model (Crawley 2007). Significant effects in the model were analysed with Tukey's contrast *post hoc* tests, performed using the package Multcomp 1.2–8 (Hothorn et al. 2008).

In the assessment of a predator's functional response, there is a range of models and choice is based on whether a particular study takes a mechanistic or phenomenological approach (Jeschke et al. 2002). Although mechanistic application of parameters such as attack rate and handling time should be supported with empirical measurements of such estimates (Caldow and Furness 2001; Jeschke and Hohberg 2008), the phenomenological use of these parameters provides a tool to examine differences in functional response Types and magnitudes in comparative experiments, as is the approach taken here (Alexander et al. 2013; Dick et al. 2013, 2014; MacNeil et al. 2013).

We first determined the functional response Type using logistic regression, testing for a negative linear coefficient (fitted using maximum likelihood) in the relationship between the proportion of prey eaten and prey density that indicates a Type II response (Trexler et al. 1988; Juliano 2001). Further, we estimated values of 'a' (attack rate), 'h' (handling time) and maximum feeding rate ( $1/hT$ , where  $T$ =experimental period) using the 'random predator equation' (Rogers 1972), which is

appropriate where prey are not replaced as they are consumed (Juliano 2001);

$$N_e = N_0 \{1 - \exp[a(N_e h - T)]\}$$


where  $N_e$  is the number of prey eaten,  $N_0$  is the initial density of prey,  $a$  is the attack constant,  $h$  is the handling time and  $T$  is the experimental period. Due to the implicit nature of the random predator equation, the Lambert W function was implemented to fit the model to the data (Bolker 2008). Bootstrapping was used to generate multiple estimates ( $n=30$ ) of the response parameters of attack rate  $a$ , handling time  $h$  and maximum feeding rate ( $1/hT$ ), which were then compared between habitat complexities in a GLM with Tukey's contrast *post hoc* tests.

## Results

In control groups with no predators, prey survival was always >98 % in each of the habitat treatments. Experimental deaths were therefore attributed to predation by juvenile largemouth bass. As habitat became less complex, significantly more prey were eaten ( $F_{3, 68}=8.41$ ,  $p<0.001$ ; Fig. 1), and bass in the highest habitat complexity consumed significantly less prey compared to all other habitat complexities (Tukey's contrasts,  $p<0.01$ ; Fig. 1). There were no differences in prey consumed between intermediate, low and zero habitat complexities. Significantly more prey were consumed at higher as compared to lower densities ( $F_{5, 68}=76.88$ ,  $p<0.001$ ).

Logistic regression indicated that, in each of the habitat complexities, largemouth bass exhibited a Type II functional response towards prey, as revealed by significantly negative linear coefficients (Table 1; Fig. 2a–d). As habitat became less complex, attack rate  $a$  significantly declined ( $F_{3, 116}=26.28$ ,  $p<0.001$ ; Fig. 3a). In comparison to low and intermediate habitats, which did not differ from each other, attack rate was significantly reduced in zero habitat treatments, and significantly greater in high complexity treatments (Tukey's contrasts, all  $p<0.01$ ; Fig. 3a). Again as habitat became less complex, handling time  $h$  declined ( $F_{3, 116}=151.12$ ,  $p<0.001$ ), and was greatest in high complexity in comparison to the other treatments (Tukey's contrasts, all  $p<0.01$ ; Fig. 3b). This was also reflected in an increase in maximum feeding rate ( $F_{3, 116}=99.09$ ,  $p<0.001$ ), that was greatest for zero and low habitat

**Fig. 1** Mean prey consumed ( $\pm$ SE) by juvenile largemouth bass in high, intermediate, low and zero complexity simulated habitats. Different letters above bars indicate significant differences (Tukey's contrasts,  $p<0.01$ )



complexities, and lowest in high complexity treatments (Tukey's contrasts, all  $p<0.01$ ; Fig. 3c).

## Discussion


Freshwater systems are threatened by a number of drivers of global change (Sala et al. 2000; Buisson et al. 2013) and, around the world, formerly pristine headwater stream environments are impacted by both habitat destruction and invasions by non-native species (Impson et al. 2002; Ellender et al. 2011). Such drivers have important consequences when considered individually, however, they may also act in combination to result in greater, synergistic impacts to native

populations (Didham et al. 2007). Furthermore, there is a pressing requirement to predict such impacts of both established and emerging invasive species under a range of environmental conditions such that appropriate mitigation and control measures may be implemented (Byers et al. 2002; Simberloff et al. 2013; Alexander et al. 2014; Dick et al. 2014).

Reduction in habitat complexity significantly increased consumption rates by invasive juvenile largemouth bass *Micropterus salmoides* of the female guppy *Poecilia reticulata*. We have thus demonstrated that the impact by juvenile largemouth bass, one of the “100 World's Worst” invaders (ISSG 2013), may be heightened with degradation of habitat. In addition, we found that between zero to intermediate structural

**Table 1** Parameter estimates (and significance levels) from logistic regression analyses of proportion of prey consumed against initial prey density, in high, intermediate, low and zero habitat complexities. Values for the intercept and linear ( $N_0$ ) terms are presented with p values

Habitat complexity	Intercept (p value)	$N_0$ (p value)	Functional response type
High	0.21 ( $p=0.39$ )	-0.04 ( $p<0.001$ )	II
Intermediate	2.04 ( $p<0.001$ )	-0.07 ( $p<0.001$ )	II
Low	1.73 ( $p<0.001$ )	-0.05 ( $p<0.001$ )	II
Zero	0.73 ( $p<0.01$ )	-0.04 ( $p<0.001$ )	II



**Fig. 2** Functional responses of juvenile bass towards prey in **a** high, **b** intermediate, **c** low and **d** zero habitat complexity (as modelled by the Rogers random predator equation for a Type II

response). Data are mean number of prey consumed at each density  $\pm$  SE


complexities, there was no significant difference in the numbers of prey consumed. This suggests the occurrence of a threshold in complexity between the intermediate and high-complexity experimental habitats that can reduce the efficiency of the predator (Coull and Wells 1983; Gotceitas and Colgan 1989; Manatunge et al. 2000). This may in turn have important consequences for predatory behaviours in instances where predators alter their foraging modes in response to changes in their surrounding environment (Scharf et al. 2006). There may also be additional consequences for predator-prey dynamics due to effects on predator-predator interactions, potentially influencing facilitation or interference outcomes (Sih et al. 1998; Warfe and Barmuta 2004).

In each of the four levels of habitat complexity, juvenile bass exhibited a Type II functional response towards the fish prey. This is counter to a number of studies demonstrating how variations in environment, such as habitat complexity, light intensity and temperature (Eggleston 1990; Koski and Johnson 2002; Alexander et al. 2012), may result in changes towards Type III responses. Generally changes in functional response types occur when factors, such as

environmental conditions, affect the searching ability of a predator. These are generally most influential at low prey densities (Crowder and Cooper 1982; Heck and Crowder 1991) and habitat complexity is often an important determinant of such outcomes (Buckel and Stoner 2000; Kushner and Hovel 2006; Alexander et al. 2012). Such a change in functional response can be significant when considering population stability and viability, as Type II responses can drive prey populations to local extinction if prey are unable to match predator consumption rate, with, for example, reproductive output (Twardochleb et al. 2012; MacNeil et al. 2013).

Although Type II responses were observed in each habitat, there were differences recorded in model parameters. Attack rates were greatest in the most complex habitat treatment and lowest when no habitat was present. As the scaling parameter of the curve, the attack rate describes the slope of the line at the lowest prey densities and therefore provides an indication of predator efficiency at these densities (Hassell and May 1973; Jeschke et al. 2002). The observed attack rates thus reflect the behaviour of a species that is predominantly an ambush predator that seeks out structure (Savino and





**Fig. 3** Mean (+SE) **a** attack rate  $a$ , **b** handling time  $h$  and **c** maximum feeding rate  $1/hT$  derived from bootstrapping ( $n=30$  each) of juvenile largemouth bass consuming prey in high,

intermediate, low and zero complexity simulated habitats. Different letters indicated significant differences (Tukey's contrasts,  $p<0.01$ )

Stein 1982, 1989b). Juvenile bass in particular may be efficient predators in dense vegetation, with smaller body size permitting comparably easier access to prey than older, larger individuals (Almany 2004b). Thus, the reduction in attack parameter in less complex habitats in comparison to the denser structure in the present study suggests that at low densities, degradation of habitat may in fact provide prey with a reduced mortality risk in comparison to more complex habitats.

As a further reflection of the greater predatory efficiency of juvenile bass at higher prey densities, differences in mean handling times, and thus maximum feeding rates, indicated greater predation at higher prey densities when habitat complexity was reduced. In comparison, high complexity structure reduced maximum feeding rates, suggesting that, although efficient predators at lower densities in these habitats, juvenile bass are impeded overall by

such structure. At zero and low habitat complexities, bass were comparably more efficient as indicated by significantly greater feeding rates, therefore with reductions in habitat cover, certain densities of prey are more vulnerable to predation by this species. This may result from the reduction in the physical barrier the habitat complexity provides with simplification in structure (Warfe and Barmuta 2004), or indeed be a consequence of reduced refuge space for prey whereby safe areas become saturated and prey are pushed out into the open where they are more vulnerable to predation (Forrester and Steele 2004).

Type II functional responses can, under certain conditions, be de-stabilising to prey populations and reduce their viability, and indeed at low prey densities in high habitat complexity areas prey populations may be driven to local extinction by juvenile bass as suggested by

the elevated attack rates. Prey populations under such circumstances may, however, be stabilised with the presence of alternative prey whereby as one species becomes rare, the predator switches to feed on another, resulting in a Type III functional response (Akre and Johnson 1979; Elliott 2004). However, field studies consistently suggest that prey populations are heavily impacted by largemouth bass (Godinho and Ferreira 2000; Almeida et al. 2012; Weyl et al. 2013), and we therefore assume that the strength of the Type II responses observed here drives lack of coexistence between bass and native prey as is congruent with other functional response studies that link experimental findings with observed field patterns (Bollache et al. 2008; Dick et al. 2013; MacNeil et al. 2013).

The use of functional responses in a predictive capacity, as applied here, to investigate what may be expected with changes to the environment when important drivers of global change interact, is a further demonstration of the utilisation of this methodology (see Dick et al. 2013, 2014). In this study, the combination of habitat degradation, as simulated with reductions in the density of a plant mimic, and the *per capita* impact of predation by invasive largemouth bass as determined by functional responses, indicates that impacts of the invasive species may potentially be greater with reductions in habitat. Largemouth bass, however, are well established in a number of systems worldwide (Welcomme 1992), and where their removal is not possible, a potential mitigation measure is therefore to focus efforts on the protection of natural vegetation and riparian zones. For further investigation it is suggested that other determinants of invader ecological impacts, such as the numerical response to examine the reproductive and/or aggregative response to prey, are quantified; however, the use of functional responses continues to be a rapid, reliable and in particular predictive assessment of the potential ecological impacts of invasive species in a changing world.

**Acknowledgements** We thank K. van der Walt, N. Songongo, G. Taylor, A. Luger, B. Ellender, D. Howell, C. Hoek, Y. Omardien, M. Pegg, B. Dredge, L. Mofu, P. Weyl and O. Weyl for technical and logistical support, including field collections. This research was funded by the DST-NRF Centre of Excellence for Invasion Biology. DMR received support from the National Research Foundation (grant 85,417). The Department of Economic Development, Environmental Affairs and Tourism (Cacadu Region) is thanked for issuing research permits (4/13CR & 5/13CR). JD acknowledges support from NERC and the Leverhulme Trust.

## References

- Akre BG, Johnson DM (1979) Switching and sigmoid functional response curves by damselfly naiads with alternative prey available. *J Anim Ecol* 48:703–720. doi:[10.2307/4191](https://doi.org/10.2307/4191)
- Alexander ME, Dick JTA, O'Connor NE et al (2012) Functional responses of the intertidal amphipod *Echinogammarus marinus*: effects of prey supply, model selection and habitat complexity. *Mar Ecol Prog Ser* 468:191–202. doi:[10.3354/meps09978](https://doi.org/10.3354/meps09978)
- Alexander ME, Dick JTA, O'Connor NE (2013) Trait-mediated indirect interactions in a marine intertidal system as quantified by functional responses. *Oikos* 122:1521–1531. doi:[10.1111/j.1600-0706.2013.00472.x](https://doi.org/10.1111/j.1600-0706.2013.00472.x)
- Alexander ME, Dick JTA, Weyl OLF et al (2014) Existing and emerging high impact invasive species are characterized by higher functional responses than natives. *Biol Lett* 10:2–6
- Almany GR (2004a) Does increased habitat complexity reduce predation and competition in coral reef fish assemblages? *Oikos* 106:275–284
- Almany GR (2004b) Differential effects of habitat complexity, predators and competitors on abundance of juvenile and adult coral reef fishes. *Oecologia* 141:105–13. doi:[10.1007/s00442-004-1617-0](https://doi.org/10.1007/s00442-004-1617-0)
- Almeida D, Almodóvar A, Nicola GG et al (2012) Trophic plasticity of invasive juvenile largemouth bass *Micropterus salmoides* in Iberian streams. *Fish Res* 113:153–158. doi:[10.1016/j.fishres.2011.11.002](https://doi.org/10.1016/j.fishres.2011.11.002)
- Anderson TW (2001) Predator responses, prey refuges, and density-dependent mortality of a marine fish. *Ecology* 82: 245–257. doi:[10.2307/2680100](https://doi.org/10.2307/2680100)
- Beukers JS, Jones GP (1997) Habitat complexity modifies the impact of piscivores on a coral reef fish population. *Oecologia* 114:50–59. doi:[10.1007/s004420050419](https://doi.org/10.1007/s004420050419)
- Bolker BM (2008) emdbook: Ecological models and data in R, R Package. Princeton University Press, Princeton, USA
- Bollache L, Dick JTA, Farnsworth KD, Montgomery WI (2008) Comparison of the functional responses of invasive and native amphipods. *Biol Lett* 4:166–169. doi:[10.1098/rsbl.2007.0554](https://doi.org/10.1098/rsbl.2007.0554)
- Boström C, Mattila J, Christo JM (1999) The relative importance of food and shelter for seagrass-associated invertebrates: a latitudinal comparison of habitat choice by isopod grazers. *Oecologia* 120:162–170. doi:[10.1007/s004420050845](https://doi.org/10.1007/s004420050845)
- Buckel JA, Stoner AW (2000) Functional response and switching behavior of young-of-the-year piscivorous bluefish. *J Exp Mar Bio Ecol* 245:25–41
- Buisson L, Grenouillet G, Villéger S et al (2013) Toward a loss of functional diversity in stream fish assemblages under climate change. *Glob Chang Biol* 19:387–400. doi:[10.1111/gcb.12056](https://doi.org/10.1111/gcb.12056)
- Bunn SE, Arthington AH (2002) Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity. *Environ Manage* 30:492–507. doi:[10.1007/s00267-002-2737-0](https://doi.org/10.1007/s00267-002-2737-0)
- Byers JE, Reichard S, Randall JM et al (2002) Directing research to reduce the impacts of nonindigenous species. *Conserv Biol* 16:630–640. doi:[10.1046/j.1523-1739.2002.01057.x](https://doi.org/10.1046/j.1523-1739.2002.01057.x)
- Caldow RWG, Furness RW (2001) Does Holling's disc equation explain the functional response of a kelp parasite? *J Anim Ecol* 70:650–662



- Coull BC, Wells JBJ (1983) Refuges from fish predation : experiments with phytal meiofauna from the New Zealand rocky intertidal. *Ecology* 64:1599–1609
- Crawley MJ (2007) *The R Book*. John Wiley, Chichester
- Crooks JA (1998) Habitat alteration and community-level effects of an exotic mussel, *Musculista senhousia*. *Mar Ecol Prog Ser* 162:137–152. doi:[10.3354/meps162137](https://doi.org/10.3354/meps162137)
- Crowder LB, Cooper WE (1982) Habitat structural complexity and the interaction between bluegills and their prey. *Ecology* 63:1802–1813
- Dick JTA, Gallagher K, Avlilas S et al (2013) Ecological impacts of an invasive predator explained and predicted by comparative functional responses. *Biol Invasions* 15:837–845
- Dick JTA, Alexander ME, Jeschke JM et al (2014) Advancing impact prediction and hypothesis testing in invasion ecology using a comparative functional response approach. *Biol Invasions* 16: 735–753
- Didham RK, Tylianakis JM, Gemmell NJ et al (2007) Interactive effects of habitat modification and species invasion on native species decline. *Trends Ecol Evol* 22:489–96. doi:[10.1016/j.tree.2007.07.001](https://doi.org/10.1016/j.tree.2007.07.001)
- Dudgeon D, Arthington AH, Gessner MO et al (2006) Freshwater biodiversity: importance, threats, status and conservation challenges. *Biol Rev* 81:163–182. doi:[10.1017/S1464793105006950](https://doi.org/10.1017/S1464793105006950)
- Eggleston DB (1990) Behavioural mechanisms underlying variable functional responses of blue crabs, *Callinectes sapidus*, feeding on juvenile oysters, *Crassostrea virginica*. *J Anim Ecol* 59:615–630. doi:[10.2307/4884](https://doi.org/10.2307/4884)
- Ellender BR, Weyl OLF, Swartz ER (2011) Invasion of a headwater stream by non-native fishes in the Swartkops river system, South Africa. *Afr Zool* 46:39–46
- Elliott JM (2004) Prey switching in four species of carnivorous stoneflies. *Freshw Biol* 49:709–720. doi:[10.1111/j.1365-2427.2004.01222.x](https://doi.org/10.1111/j.1365-2427.2004.01222.x)
- Facon B, Genton BJ, Shykoff J et al (2006) A general eco-evolutionary framework for understanding bioinvasions. *Trends Ecol Evol* 21:130–135. doi:[10.1016/j.tree.2005.10.012](https://doi.org/10.1016/j.tree.2005.10.012)
- Fazey I, Fischer J, Lindenmayer DB (2005) What do conservation biologists publish? *Biol Conserv* 124:63–73. doi:[10.1016/j.biocon.2005.01.013](https://doi.org/10.1016/j.biocon.2005.01.013)
- Flynn AJ, Ritz DA (1999) Effect of habitat complexity and predatory style on the capture success of fish feeding on aggregated prey. *J Mar Biol Assoc United Kingdom* 79:487–494
- Forrester GE, Steele MA (2004) Predators, prey refuges and the spatial scaling of density-dependent prey mortality. *Ecology* 85:1332–1342
- Godinho FN, Ferreira MT (2000) Composition of endemic fish assemblages in relation to exotic species and river regulation in a temperate stream. *Biol Invasions* 2:231–244
- Gosnell JS, DiPrima JB, Gaines SD (2012) Habitat complexity impacts persistence and species interactions in an intertidal whelk. *Mar Biol* 159:2867–2874. doi:[10.1007/s00227-012-2047-x](https://doi.org/10.1007/s00227-012-2047-x)
- Gotceitas V, Colgan P (1989) Predator foraging success and habitat complexity: quantitative test of the threshold hypothesis. *Oecologia* 80:158–166
- Griffen BD, Delaney DG (2007) Species invasion shifts the importance of predator dependence. *Ecology* 88:3012–3021
- Hassell MP (1978) Functional responses. In: Hassell M (ed) *Dyn. Arthropod Predator-Prey Syst.* Princeton University Press, Princeton, pp 28–49
- Hassell MP, May RM (1973) Stability in insect Host-parasite models. *J Anim Ecol* 42:693–726
- Heck KL, Crowder LB (1991) Habitat structure and predator-prey interactions in vegetated aquatic systems. In: Bell S, McCoy E, Mushinsky H (eds) *Habitat struct. physical arrange. objects Sp.* Chapman and Hall, London, pp 281–299
- Holling C (1959) Some characteristics of simple types of predation and parasitism. *Can Entomol* 91:385–398
- Hothorn T, Bretz F, Westfall P (2008) Simultaneous inference in general parametric models. *Biometrical J* 50:346–363
- Impson ND, Bills IR, Cambray JA (2002) A conservation plan for the unique and highly threatened freshwater fishes of the Cape Floral Kingdom. In: Collares-Pereira MJ, Coelho MM, Cowx I (eds) *Conserv. Freshw. Fishes Options Futur.* Blackwell Science, Oxford, pp 432–440
- Jeschke JM, Hohenberg K (2008) Predicting and testing functional responses: an example from a tardigrade–nematode system. *Basic Appl Ecol* 9:145–151. doi:[10.1016/j.baec.2007.01.006](https://doi.org/10.1016/j.baec.2007.01.006)
- Jeschke JM, Kopp M, Tollrian R (2002) Predator functional response: discriminating between handling and digesting prey. *Ecol Monogr* 72:95–112. doi:[10.2307/3100087](https://doi.org/10.2307/3100087)
- Jones CLW (2003) Intercohort cannibalism and parturition-associated behaviour of captive-bred swordtail, *Xiphophorus helleri* (Pisces: Poeciliidae), PhD thesis, Rhodes University, Grahamstown, 215 pp.
- Juliano SA (2001) Nonlinear curve fitting: predation and functional response curves. In: Scheiner SM, Gurevitch J (eds) *Des. anal. ecol. exp*, 2nd ed. Oxford University Press, Oxford, pp 178–196
- Koski ML, Johnson BM (2002) Functional response of kokanee salmon (*Oncorhynchus nerka*) to Daphnia at different light levels. *Can J Fish Aquat Sci* 59:707–716
- Kushner RB, Hovel KA (2006) Effects of native predators and eelgrass habitat structure on the introduced Asian mussel *Musculista senhousia* (Benson in Cantor) in southern California. *J Exp Mar Bio Ecol* 332:166–177
- Levine JM, D'Antonio CM (2003) Forecasting biological invasions with increasing international trade. *Conserv Biol* 17: 322–326. doi:[10.1046/j.1523-1739.2003.02038.x](https://doi.org/10.1046/j.1523-1739.2003.02038.x)
- Lipcius RN, Hines AH (1986) Variable functional responses of a marine predator in dissimilar homogeneous microhabitats. *Ecology* 67:1361–1371
- MacNeil C, Dick JTA, Alexandre ME et al (2013) Predators vs. alien: differential biotic resistance to an invasive species by two resident predators. *NeoBiota* 19:1–19. doi:[10.3897/neobiota](https://doi.org/10.3897/neobiota)
- Manatunge J, Asaeda T, Priyadarshana T (2000) The influence of structural complexity on fish-zooplankton interactions: a study using artificial submerged macrophytes. *Environ Biol Fishes* 58:425–438
- Mokany K, Harwood TD, Williams KJ, Ferrier S (2012) Dynamic macroecology and the future for biodiversity. *Glob Chang Biol* 18:3149–3159. doi:[10.1111/j.1365-2486.2012.02760.x](https://doi.org/10.1111/j.1365-2486.2012.02760.x)
- Murdoch WW, Oaten A (1975) Predation and population stability. *Adv Ecol Res* 9:1–131
- Nelson WG, Bonsdorff E (1990) Fish predation and habitat complexity : are complexity thresholds real ? *J Exp Mar Bio Ecol* 141:183–194
- Nilsson C, Berggren K (2000) Alterations of riparian ecosystems caused by river regulation. *Bioscience* 50:783–792
- Olson MH, Young BP, Blinkoff KD (2003) Mechanisms underlying habitat use of juvenile largemouth bass and smallmouth

- Bass. *Trans Am Fish Soc* 132:37–41. doi:[10.1577/1548-8659\(2003\)132<0398](https://doi.org/10.1577/1548-8659(2003)132<0398)
- Parker IM, Simberloff D, Lonsdale WM et al (1999) Impact : toward a framework for understanding the ecological effects of invaders. *Biol Invasions* 1:3–19
- Persson L, Eklöv P (1995) Prey refuges affecting interactions between piscivorous perch and juvenile perch and roach. *Ecology* 76:70–81
- Radomski P, Goeman TJ (2001) Consequences of human lake-shore development on emergent and floating-leaf vegetation abundance. *North Am J Fish Manag* 21:46–61
- Ricciardi A, Hoopes MF, Marchetti MP, Lockwood JL (2013) Progress toward understanding the ecological impacts of nonnative species. *Ecol Monogr* 83:263–282
- Rogers D (1972) Random search and insect population models. *J Anim Ecol* 41:369–383. doi:[10.2307/3474](https://doi.org/10.2307/3474)
- Rowles AD, O'Dowd DJ (2006) Interference competition by Argentine ants displaces native ants: implications for biotic resistance to invasion. *Biol Invasions* 9:73–85. doi:[10.1007/s10530-006-9009-5](https://doi.org/10.1007/s10530-006-9009-5)
- Saha N, Aditya G, Saha GK (2009) Habitat complexity reduces prey vulnerability: an experimental analysis using aquatic insect predators and immature dipteran prey. *J Asia Pac Entomol* 12:233–239. doi:[10.1016/j.aspen.2009.06.005](https://doi.org/10.1016/j.aspen.2009.06.005)
- Sala OE, Stuart Chapin F III et al (2000) Global biodiversity scenarios for the year 2100. *Science* 287(80):1770–1774. doi:[10.1126/science.287.5459.1770](https://doi.org/10.1126/science.287.5459.1770)
- Saunders DL, Meeuwig JJ, Vincent ACJ (2002) Freshwater protected areas: strategies for conservation. *Conserv Biol* 16:30–41. doi:[10.1046/j.1523-1739.2002.99562.x](https://doi.org/10.1046/j.1523-1739.2002.99562.x)
- Savino JF, Stein RA (1982) Predator-prey interaction between largemouth bass and bluegills as influenced by simulated, submersed vegetation. *Trans Am Fish Soc* 111:255–266. doi:[10.1577/1548-8659\(1982\)111<255](https://doi.org/10.1577/1548-8659(1982)111<255)
- Savino JF, Stein RA (1989a) Behavior of fish predators and their prey - habitat choice between open water and dense vegetation. *Environ Biol Fish* 24:287–293
- Savino JF, Stein RA (1989b) Behavioural interactions between fish predators and their prey: effects of plant density. *Anim Behav* 37:311–321
- Scharf I, Nulman E, Ovadia O, Bouskila A (2006) Efficiency evaluation of two competing foraging modes under different conditions. *Am Nat* 168:350–357. doi:[10.1086/506921](https://doi.org/10.1086/506921)
- Sih A, Englund G, Wooster D (1998) Emergent impacts of multiple predators on prey. *Trends Ecol Evol* 13:350–355
- Simberloff D, Martin J-L, Genovesi P et al (2013) Impacts of biological invasions: what's what and the way forward. *Trends Ecol Evol* 28:58–66. doi:[10.1016/j.tree.2012.07.013](https://doi.org/10.1016/j.tree.2012.07.013)
- Solomon ME (1949) The natural control of animal populations. *J Anim Ecol* 18:1–35
- Thomas CD, Cameron A, Green RE et al (2004) Extinction risk from climate change. *Nature* 427:145–148. doi:[10.1038/nature02121](https://doi.org/10.1038/nature02121)
- Trexler JC, McCulloch CE, Travis J (1988) How can the functional response best be determined. *Oecologia* 76:206–214
- Twadodchleb LA, Novak M, Moore JW (2012) Using the functional response of a consumer to predict biotic resistance to invasive prey. *Ecol Appl* 22:1162–1171
- Warfe DM, Barmuta LA (2004) Habitat structural complexity mediates the foraging success of multiple predator species. *Oecologia* 141:171–178
- Welcomme RL (1992) A history of international introductions of inland aquatic species. *ICES J Mar Sci* 184:3–14
- Weyl OLF, Ellender BR, Woodford DJ, Jordaan MS (2013) Fish distributions in the Rondegat River, Cape Floristic Region, South Africa, and the immediate impact of rotenone treatment in an invaded reach. *African J Aquat Sci* 1–9. doi:[10.2989/16085914.2012.753401](https://doi.org/10.2989/16085914.2012.753401)